

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Anthocyanins in Berries and Their Potential Use in Human Health

---

Daniela Peña-Sanhueza,  
Claudio Inostroza-Blancheteau,  
Alejandra Ribera-Fonseca and Marjorie Reyes-Díaz

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67104>

---

## Abstract

Anthocyanin pigments are responsible for the red, purple, and blue colors of many fruits, vegetables, cereal grains, and flowers, increasing the interest due to their strong antioxidant capacity and their possible use to the benefit of human health. Abundant evidence is available about the preventive and therapeutic roles of anthocyanin in different kinds of chronic diseases. According to the structural differences and anthocyanin content of berries such as blackberry, blueberry, chokeberry, and others, there are different healthy properties in the treatments of circulatory disorders, cancer cell lines, and diabetes as well as antiviral and antimicrobial activities. On the other hand, molecular aspects play an important role in anthocyanin biosynthesis, making it possible to determine how biotic and abiotic factors impact its biosynthesis complex. Thus, the aim of this chapter was to describe the use of anthocyanins from berries for human health and their potential use as a pharmacological bioresource in the prevention of chronic diseases. In addition, an update of the molecular mechanisms involved in anthocyanin biosynthesis will be discussed.

**Keywords:** anthocyanins, berries, cancer, transcription factors

---

## 1. Introduction

The scientific evidence regarding the positive relationship between diet and health has increased consumer demand for more information related to healthy diets, including fruits and vegetables, with functional characteristics that help to delay the aging process and

reduce the risk of several diseases, mainly cardiovascular diseases and cancer [1]. Berries are recognized as an important component of healthy diets due to their bioactive compounds. In this sense, commercial berry species such as blackberry (*Rubus* sp.), bilberry (*Vaccinium myrtillus* L.), blackcurrant (*Ribes rugrum* L.), chokeberry (*Aronia melanocarpa* (Michx.) Elliott.), cranberry (*V. macrocarpon* Ait.), bayberry (*Myrica* sp.), raspberry (*Rubus ideaus* L.), black raspberry (*Rubus occidentalis* L.), strawberry (*Fragaria ananassa* Duch.), highbush blueberry (*V. corymbosum* L.), maqui (*Aristotelia chilensis*), murtilla (*Ugni molinae* Turcz.), and calafate (*Berberis microphylla* G. Forst.) are particularly rich sources of antioxidants, which are usually consumed in fresh and processed products [2–5]. Higher plants, especially berry species, synthesize a diverse group of phenolic compounds such as flavonoids. These plant secondary metabolites have many biological functions, including their key role in plant-microbe interaction, plant-pathogen interaction, pollen-tube growth, UV radiation protection, tissue pigmentation, and others [6, 7]. Flavonoid compounds, which include flavonols, flavones, flavanols, flavanones, isoflavonoids, and anthocyanins, are molecules widely accumulated in vascular plants and to a lesser extent in mosses, being accumulated in all organs and tissues at different stages of development and depending on the environmental conditions [6].

Anthocyanins are natural pigments responsible for the blue, purple, red, and orange colors of many fruits and vegetables [8, 9]. Anthocyanins are a glycoside form of anthocyanidins [9], and the structural differences among them are related to the number of hydroxyl group, position, and kind and/or number of sugars linked to the molecule [10, 11]. These compounds appear to be an interesting natural resource of water-soluble dyes because they are easily incorporated in aqueous media [12]. Another important property of anthocyanins is their remarkable antioxidant activity, playing a vital role in the prevention of neuronal and cardiovascular illnesses, diabetes, cancer, etc. [11, 13]. Many reports have focused on the effect of anthocyanins in cancer prevention [14], human nutrition [15], and their biological activity [10]. Nowadays, there is an increased interest in explaining the role of anthocyanins as a natural antioxidant and their mechanism of action on human health as well as the treatment of chronic diseases and their use as a natural dye, substituting the synthetic dyes, which can be toxic to humans. This review endeavors to describe the use of anthocyanins from berries for human health and their potential use as a pharmacological bioresource in the prevention of chronic diseases. In addition, an update of the molecular mechanisms involved in anthocyanin biosynthesis will be discussed. Finally, recent clinical and preclinical studies about anthocyanin use in the prevention of human diseases are reported.

## 2. Anthocyanin and phenolic compounds in berries

Phenolic acid, organic acids, tannins, anthocyanins, and flavonoids are phenolic bioactive compounds with a high concentration in the berry fruits [16]. The chemical structure of phenolic compounds is characterized by one or more aromatic rings with hydroxyl groups. According to their structural characteristics, phenolic compounds are classified into five

major groups: phenolic acids, stilbenes, flavonoids (flavonols or catechins, flavonols, flavones, flavonones, isoflavonoids, anthocyanins), tannins, and lignans [13]. The concentration of phenolic compounds in berry fruits is altered by many factors, such as genotype, species, agronomic management, climatic factors, ripening stage, harvesting time, and postharvest management [17, 18]. Given the plant phenol attributes of berry species, attention has largely focused on anthocyanin and flavonol antioxidant action on human health. In this way, substantial epidemiological and experimental research suggests that intakes of recognized nutritional antioxidants such as vitamin E and carotenoids can decrease the oxidative damage of proteins, lipids, and DNA *in vivo* and may reduce the incidence of developing many chronic diseases in humans [19]. The *in vitro* antioxidant effectiveness of anthocyanins and other polyphenols is due to its donation of free hydrogen atom from an aromatic hydroxyl group of the antioxidant molecules, acting as radical scavenger [20].

It has been reported that the antioxidant capacity of flavonoids is stronger than vitamins C and E [21, 22], and under *in vitro* conditions, flavonoids can prevent injury in different ways, acting as a suppressor of reactive oxygen formation, scavenging free radicals by hydrogen atom donation [22, 23], activating antioxidant enzymes [23, 24], chelating metal, reducing  $\alpha$ -tocopheryl radicals, inhibiting oxidases, oxidative stress mitigation by nitric oxide, increasing uric acid levels, and increasing antioxidant properties of low-molecular antioxidants [22]. Anthocyanin concentration in blackberry is much higher than in raspberry and strawberry and similar to red currant blueberry, depending on the cultivar (see **Table 1**).

Anthocyanin concentration widely differs significantly among plant species, even among species of the same genus. In **Table 1**, anthocyanin and total phenolic compounds of different species and cultivars and their analysis are detailed. In blackberry, anthocyanin content is generally similar in all species, but phenolic content shows strong differences (**Table 1**). Anthocyanin content in *Rubus insularis* F. Aresch. represents 36% of the phenolic compounds, whereas in *R. fruticosus* cultivar Hull Thornless, it only represents 6.4% of the total phenolic compounds (**Table 1**). Raspberry (*R. innominatus* S. Moore) showed higher anthocyanin level, representing 41.2% of the total phenolic compounds. *R. ideaus* show high phenolic compounds; however, their high content does not necessarily represent a high anthocyanin content. *R. ideaus* Heritage cultivar has showed the highest anthocyanin percent with respect to the total phenolic compounds, representing 3.8% (**Table 1**). Additionally, blueberry cultivars showed low differences between anthocyanin and phenolic compounds, but they showed greater health benefits than other berries due to their particularly high proportion of anthocyanins. In some cases, high anthocyanin content in blueberries is related to high antioxidant capacity, but the anthocyanin contents and composition are different in each species and cultivar (**Table 1**). More specifically, the *V. corymbosum* cultivar (Duke) contains 63% anthocyanin with respect to the total phenolic compounds, followed by the cultivars CVAC5.001 and Brigitta, with 46 and 41%, respectively, and finally by Bluecrop with 27%. It is therefore necessary to evaluate the correlation between anthocyanin content and total phenolic compounds, because the ratio can exist between the two parameters, but it is not necessary to estimate in all species or among cultivars of the same genus (**Table 1**) [25–29].

| Scientific name                          | Common name  | Cultivar         | Anthocyanins* | Phenolics** | References |
|--|--------------|------------------|---------------|-------------|------------|
| <i>Rubus cyri</i> Juz.                   | Blackberry   | Native           | 143           | 545         | [25]       |
| <i>Rubus georgicus</i> Focke             | Blackberry   | Native           | 89            | 561         | [25]       |
| <i>Rubus insularis</i> F. Aresch.        | Blackberry   | Native           | 170           | 472         | [25]       |
| <i>Rubus ursinus</i> (Douglas ex Hook.)  | Blackberry   | Native           | 211           | 629         | [25]       |
| <i>Rubus fruticosus</i> L.               | Blackberry   | Chactaw          | 125           | 1703        | [26]       |
| <i>Rubus fruticosus</i>                  | Blackberry   | T. evergreen     | 146           | 2061        | [26]       |
| <i>Rubus fruticosus</i>                  | Blackberry   | Hull Thornless   | 152           | 2349        | [26]       |
| <i>Rubus idaeus</i> L.                   | Raspberry    | Native           | 65            | 517         | [27]       |
| <i>Rubus innominatus</i> S. Moore        | Raspberry    | Native           | 52            | 126         | [25]       |
| <i>Rubus niveus</i> Thunb.               | Raspberry    | Native           | 230           | 402         | [25]       |
| <i>Rubus ideaus</i>                      | Raspberry    | Heritage         | 49            | 1280        | [26]       |
| <i>Rubus ideaus</i>                      | Raspberry    | Autumm Bliss     | 39            | 2494        | [26]       |
| <i>Rubus ideaus</i>                      | Raspberry    | Fallgold         | 3             | 1459        | [26]       |
| <i>Rubus ideaus</i>                      | Raspberry    | Meeker           | 42            | 2116        | [26]       |
| <i>Ribes sativum</i>                     | Red currants | London Market    | 7.8           | 1115        | [26]       |
| <i>Ribes sativum</i> (Lam.) Mert. & Kock | Red currants | Rovada           | 7.5           | 1193        | [26]       |
| <i>Ribes sativum</i>                     | Red currants | White Versailles | 1.4           | 657         | [26]       |
| <i>Ribes nigrum</i> L.                   | Red currants | Alagan           | 169           | 694         | [25]       |
| <i>Ribes nigrum</i>                      | Red currants | Ben Lomond       | 261           | 933         | [25]       |
| <i>Ribes nigrum</i>                      | Red currants | Ojebyn           | 165           | 830         | [25]       |
| <i>Ribes nigrum</i>                      | Red currants | Consort          | 411           | 1342        | [25]       |
| <i>Vaccinium corymbosum</i> L.           | Blueberry    | Bluecrop         | 84            | 304         | [25]       |
| <i>Vaccinium corymbosum</i>              | Blueberry    | Briggita         | 103           | 246         | [25]       |
| <i>Vaccinium corymbosum</i>              | Blueberry    | Duke             | 173           | 274         | [25]       |
| <i>Vaccinium corymbosum</i>              | Blueberry    | CVAC5.001        | 430           | 868         | [25]       |
| <i>Vaccinium corymbosum</i>              | Blueberry    | Native           | 62-235        | 181-473     | [28]       |
| <i>Vaccinium corymbosum</i>              | Blueberry    | Bluegold         | 206           | 432         | [29]       |
| <i>Vaccinium corymbosum</i>              | Blueberry    | Briggita         | 190           | 468         | [29]       |
| <i>Vaccinium corymbosum</i>              | Blueberry    | Legacy           | 226           | 570         | [29]       |
| <i>Vaccinium angustifolium</i> Ait.      | Blueberry    | Native           | 208           | 692         | [25]       |
| <i>Vaccinium myrtillus</i> L.            | Bilberry     | Native           | 300           | 525         | [28]       |

\*mg cyanidin 3-glucoside eq./100 g fresh weight.  
\*\*mg gallic acid eq./100 g fresh weight.

**Table 1.** Total anthocyanin and phenolic content of berry fruits.

Berry species with higher anthocyanin content are interesting for use in breeding programs for increasing their content in fruits, enhancing their antioxidant capacity, and obtaining fruit products with health properties. In addition, the understanding of the molecular network of genes involved in anthocyanin biosynthesis and how biotic and abiotic factors could affect their concentration and gene regulation are a key to use it in genetic engineering and agronomic management.

### 3. Molecular regulation of anthocyanin biosynthesis

Six structural genes are common in the anthocyanin pathway in all angiosperms, which are divided into two main groups. The first group is the upstream genes or early biosynthesis genes, for example, chalcone synthase (CHS), chalcone flavanone isomerase (CHI), and flavanone 3-hydroxylase (F3H), coding for enzymes that produce precursors for one or more important non-anthocyanin flavonoids. The second group is the downstream genes or late biosynthesis genes, for example, anthocyanidin synthase (ANS), dihydroflavonol-4-reductase (DFR), and UDP-glucose flavonoid 3-oxy-glucosyltransferase (UGT), coding for enzymes specific to anthocyanin synthesis [30–32]. In the anthocyanin pathway, L-phenylalanine is converted to naringenin by phenylalanine ammonia lyase (PAL), cinnamate 4-hydroxylase (C4H), 4-coumarate CoA ligase (4CL), chalcone synthase (CHS), and chalcone isomerase (CHI). Then, the next pathway is catalyzed by the formation of complex aglycone and anthocyanin composition by flavanone 3-hydroxylase (F3H), flavonoid 3'-hydroxylase (F3'H), dihydroflavonol 4-reductase (DFR), anthocyanidin synthase (ANS), UDP-glucoside flavonoid glucosyltransferase (UGT), and methyl transferase (MT) [33]. It has been described that the transcription of early and late biosynthesis genes to produce anthocyanins appears to be regulated by R2R3-MYB and basic helix-loop-helix (bHLH, also known as MYC) called transcription factors in collaboration with tryptophan-aspartic acid repeat (WDR) or WD40 proteins [32, 34–37].

#### 3.1. MYB transcription factor

The MYB transcription factors involved in the flavonoid pathway have been identified and described for several kinds of model plants, crops, and ornamental plants. The first identified and reported MYB transcription factor in plants was in *Zea mays*, which included C1 (Colorless 1) and PL1 (Purple Leaf 1) [38]. The MYB transcription factors are composed of the so-called N-terminal MYB domain, consisting of one to three imperfect repeats of almost 52 amino acids (R1, R2, and R3), beginning with R2R3, the most abundant subfamily in plants [39]. The MYB domain is involved in DNA binding and dimerization. The C-terminal region is responsible for establishing protein-protein and regulates activation or repression of gene expression [34, 40, 41]. The MYB genes are exclusive to eukaryotic organisms [42]. In animals, these genes are associated with cell proliferation and differentiation [43, 44], whereas in plants, MYB is associated with responses to different biotic and abiotic stressors (drought, cold, pathogen disease resistance), plant development (trichome formation, seed development), stomatal movement, and many other functions [34, 40, 45, 46]. Anthocyanin biosynthesis mediated by MYB transcription factors has been reported in *Arabidopsis thaliana*



(L.) Heynh. [41, 47–49], strawberry (*F. ananassa*) [50], Chilean strawberry (*Fragaria chiloensis* (L.) Duch.) [51], apple (*Malus domestica* Borkh.) [52–54], and tomato (*Solanum lycopersicum* L.) [55]. Grape (*Vitis vinifera* L.) is the main plant species studied in this way due to its agricultural and commercial importance worldwide. Thus, many MYB transcription factors have been reported in this species by different researchers. *VvMYBPA1* and *VvMYBPA2* are involved in proanthocyanidin synthesis [46, 56], while *VvMYBF1* regulates flavonol synthesis [57]. In addition, *MYBA1* and *MYBA2* genes control the last biosynthetic step of anthocyanin synthesis [58, 59]. It is reported that a glycosylation reaction mediated by the UDP-glucose flavonoid-3-O-glucosyltransferase (UGT) enzyme produces anthocyanins in grapes [31, 39]. It is important to highlight that MYB transcription factor is conserved between different species and is one of the most important primary proteins involved in structural and biological functions. Furthermore, MYB transcription factor regulates the flavonoid pathway apparently in two ways: (a) due to variations in the C-terminal region of the protein or (b) modulating the interaction with DNA, bHLH, and WD40 protein [34, 60].

### 3.2. Basic helix-loop-helix (bHLH)

After MYB, bHLH proteins, also known as MYC, are the second most important family of transcription factors involved in anthocyanin biosynthesis [34, 61]. The bHLH protein domain is constituted of about 60 amino acids and is characterized by the presence of 19 conserved amino acids, five in the basic region, five in the first helix, one in the loop, and eight in the final second helix [61]. The basic region of bHLH has basic residues (5.8 on average) essential for DNA binding. In Arabidopsis, 20% of bHLH transcription factors do not have this domain and can act as a repressor because forming heterodimers are unable to bind to DNA [61]. Two cis-element boxes have been reported to bind with bHLH proteins, the E-box (5'-CANNTG-3'), and G-box (5'-CACGTG-3') elements. The G-box is the most commonly recognized sequence representing 81% of the proteins predicted to bind DNA [61, 62]. In the basic region, two amino acids conferred the property on binding DNA in Arabidopsis plants. The Glu13 and Arg16 are the E-box recognition motif [63]. Glu13 has contact with CA bases of E-box and Arg16, apparently helping Glu13 to bind and stabilize. In G-box, specific stabilization is mediated by His/Lys9, Glu13, and Arg17. The Arg17 interacts with inner G base, and His/Lys9 interacts with the last G of the G-box [61, 62]. The alpha-helix function is involved in homo- and hetero-dimerization and is formed by hydrophobic residues of isoleucine, leucine, and valine [34, 61]. Arabidopsis has been demonstrated that this residue is conserved in all bHLH proteins, indicating the importance of the basic region of the bHLH transcription factor in DNA binding [61, 63]. The second helix is involved in DNA binding through direct contact with the E-box. Finally, the loop is responsible for the three-dimensional arrangement of alpha-helices, and residues from the first helix loop junction are involved in association with bHLH proteins [34, 61, 63, 64]. Basic helix-loop-helix transcription factors in plants are involved in processes such as flower development [65, 66], hormonal response [67, 68], metal homeostasis [69], and others. Regarding bHLH and their relation to flavonoid synthesis, the first bHLH involved in this pathway was detected in maize in 1989 [70]. In this context, in *Z. mays* (ZmB, ZmR, and ZmLc), bHLH is involved in the regulation of the anthocyanin pathway [70–72], and

ZmIn1 is involved in the repression of flavonoid gene expression in maize aleurone [73]. In *A. thaliana*, it has been reported that *AtTT8* gene encodes a bHLH transcription factor involved in the control of proanthocyanidins and anthocyanins in seeds and seedlings [74]. Quatroccio et al. [75, 76]. reported PhAN1, PhJAF13 hBLH transcription factor from *Petunia hybrida* as being involved in the control of the anthocyanin pathway in flowers. For *Vitis vinifera*, VvMYCA1 (also known as bHLH) was reported as involved in promotion of anthocyanin accumulation in grape cells [37].

### 3.3. WDR proteins

Tryptophan-aspartic acid repeat protein (WDR) or WD40 proteins are characterized by around 44–66 amino acids, delimited by the GH dipeptide on the N-terminal size (11–24 residues from the N-terminus) and the WD dipeptide on the C-terminus [34, 77]. In Arabidopsis, WDR protein contains four (or more) tandem repeats composed of around 40 amino acids [78]. In contrast to the majority of proteins, WDR is not involved in catalytic activities such as DNA binding or gene expression regulation, mostly acting as a platform due to its capacity to interact with more than one protein at the same time [34, 78]. The work of WDR involves eukaryotic cellular process such as cell division, vesicle formation, signal transduction, RNA processing, and transcription regulation [78]. On the other hand, MYB and bHLH transcription factors have few WDR proteins involved in the flavonoid pathway, as shown in *Z. mays* (ZmPAC1), where it regulates the anthocyanin pathway in seed aleurone [79]. In Arabidopsis (*AtTTG1*), WDR proteins control trichomes, root hair, and seed mucilage production [80]. In petunia, AN11 regulates anthocyanin production as well as the pH of the flower vacuole [81], whereas in grape, *V. vinifera* WDR1 contributes to anthocyanin accumulation [37]. Although WDR proteins are not directly involved in the flavonoid pathway, particularly in anthocyanin synthesis, it is important to note that these proteins are highly conserved among species [34]. Nevertheless, few WDR proteins have been reported in plants, and it must be highlighted that WDR is involved in several metabolic and physiological processes [79, 80, 82]. To clarify the characteristics of WDR proteins and the complex formed with MYB and bHLH, which is involved in anthocyanin biosynthesis, species such as petunia and Arabidopsis have been used [34, 35].

### 3.4. MYB-bHLH-WDR (MBW complex)

MBW complex has been reported in Arabidopsis, petunia, and some varieties of grape [35, 82]. The most important function of these transcription factors is involved in the process related to DNA binding, activation of gene expression involved in the flavonoid pathway, and stabilization of the three-dimensional configuration of the complex [34]. Basic helix-loop-helix-WDR interaction is needed to WDR protein translocation into the nucleus, and this was demonstrated in onion cells using green protein fluorescent (GPF), which when expressed alone is localized in the cytosol, whereas its co-expression with PFW and MYC-RP enables the transport and localization in the nucleus [35]. The AN11 from petunia showed the same results, being detected in the cytosol [81]. *V. vinifera* subjected to high salt concentrations showed a cultivar-dependent response for anthocyanin accumulation, which was correlated with the expression of MYBA1-2, MYCA1 and WDR1 genes [37].



#### 4. Antioxidant capacity of anthocyanins in berries and their use in human health

The radical scavenging activity (RSA) of anthocyanins is largely due to the presence of hydroxyl groups in position 3 of ring C and also in the 3', 4', and 5' positions in ring B of the molecule. In general, RSA of anthocyanidins (aglycons) is superior to their respective anthocyanins (glycosides), and this decreases when the number of sugar increases [16]. Hanachi et al. [83] showed that fruits of *Berberis vulgaris* L. (barberry) have a high antioxidant activity, reducing the viability of cell cultures associated with liver cancer (HepG2). Furthermore, extracts of leaves and twigs of *B. vulgaris* have more antioxidants than fruits. Končić et al. [84] studied the antioxidant activity of extracts of leaves, branches, and roots of two species of *B. vulgaris* and *Berberis croatica* and demonstrated that all these organs exhibited antioxidant activity. In all cases, the activity was positively correlated with the content of phenolic acids and flavonols, and the flavonols played the main role in the total antioxidant activity of the studied species [84]. They also concluded that the antioxidant activities were significantly different (being higher in *B. croatica* than *B. vulgaris*) and among organs (being higher in leaves followed by branches and roots). The result of the anthocyanin concentration in different organs besides the fruits is interesting, because acquisition of anthocyanin in every season of the year has advantages for making new products with health properties. Thus, interesting results such as a new natural resource for promoting these compounds for human health have been reported. Končić et al. [84] suggested that studies into different species are needed to analyze all the organs of the plant, not just the fruits. Shin et al. [85] reported that in human liver cancer HepG2, cell proliferation was inhibited by strawberry extracts. Moreover, Chang et al. [86] reported that *Hibiscus sabdariffa* Linne (roselle) anthocyanin extracts mediated the apoptosis of human promyelocytic leukemia cells via the p38/Fas and Bid pathways. Research examining the use of black currant extract (BCE) with high concentrations of phenolic compounds on antiproliferative activity against gastric cancer SGC-7901 cells showed a positive antiradical activity and anticarcinogenic effects [87]. Moreover, extracts of mulberry showed an inhibition on the growth of human gastric carcinoma cells [88]. In this study, anthocyanins extracted from mulberry had notable promotive effects on the p38/jun/Fas/FasL and p38/p53/Bax signaling pathways, which accounted for its *in vitro* and *in vivo* growth-inhibitory and apoptotic responses in AGS (gastric cancer) cells. The effects of berries on diseases are shown in **Table 2**.

With respect to *in vivo* studies, Wang and Stoner [89] reported the effect of an anthocyanin-rich extract from black raspberries on the development of tumors in rat esophagus by N-nitrosomethylbenzylamine (NMBA), the most potent inducer of tumors in rat esophagus. This extract inhibited cell proliferation, inflammation, and induced apoptosis in the esophageal tissues (**Table 2**). Stoner et al. [90] compared the effect of black raspberry, red raspberry, strawberry, and blueberry anthocyanin and ellagitannins in fruit extract on the prevention of esophageal cancer induced by N-nitrosomethylbenzylamine (NMBA) in rats. Inhibition of NMBA-induced tumorigenesis in the rat esophagus was observed. The authors detected a reduction in cytokine levels in serum, interleukin 5 (IL-5), and GRO/KC, which is the rat homolog for human interleukin-8 (IL-8), and these cytokines

| Disease          | Scientific name                  | Common name       | Compound                       | Experimental conditions | Reference |
|------------------|----------------------------------|-------------------|--------------------------------|-------------------------|-----------|
| Liver cancer     | <i>Fragaria x ananassa</i> Duch. | Strawberry        | Crude extract                  | <i>In vitro</i>         | [85]      |
| Leukemia         | <i>Hibiscus sabdariffa</i> L.    | Rosselle          | Anthocyanin rich extract       | <i>In vitro</i>         | [86]      |
| Gastric cancer   | <i>Morus alba</i> L.             | Mulberry          | Anthocyanins                   | <i>In vitro</i>         | [88]      |
| Gastric cancer   | <i>Ribes nigrum</i> L.           | Black currant     | Crude extract                  | <i>In vitro</i>         | [87]      |
| Colon cancer     | <i>Vaccinium myrtillus</i> L.    | Bilberry          | Anthocyanin-rich extract       | <i>In vivo</i> (rats)   | [91]      |
| Colon cancer     | <i>Aronia melanocarpa</i> E.     | Chokeberry        | Anthocyanin-rich extract       | <i>In vivo</i> (rats)   | [91]      |
| Colon cancer     | <i>Vitis vinifera</i> L.         | Grape             | Anthocyanin-rich extract       | <i>In vivo</i> (rats)   | [91]      |
| Esophagus cancer | <i>Rubus occidentalis</i> L.     | Black raspberries | Anthocyanin-rich extract       | <i>In vivo</i> (rats)   | [89]      |
| Esophagus cancer | <i>Rubus occidentalis</i>        | Black raspberries | Anthocyanins and ellagitannins | <i>In vivo</i> (rats)   | [90]      |
| Esophagus cancer | <i>Rubus ideaus</i> L.           | Red raspberries   | Anthocyanins and ellagitannins | <i>In vivo</i> (rats)   | [90]      |
| Esophagus cancer | <i>Fragaria ananassa</i>         | Strawberries      | Anthocyanins and ellagitannins | <i>In vivo</i> (rats)   | [90]      |
| Esophagus cancer | <i>Vaccinium corymbosum</i> L.   | Blueberries       | Anthocyanins and ellagitannins | <i>In vivo</i> (rats)   | [90]      |
| Hepatic cancer   | <i>Berberis vulgaris</i> Duch.   | Barberries        | Crude extract                  | <i>In vivo</i> (rats)   | [83]      |
| Liver cancer     | <i>Vaccinium corymbosum</i>      | Blueberries       | Anthocyanin extract            | <i>In vitro</i> (mice)  | [94]      |
| Liver cancer     | <i>Berberis vulgaris</i>         | Barberries        | Crude extract                  | <i>In vitro</i>         | [83]      |
| Oral cancer      | <i>Rubus occidentalis</i>        | Black raspberries | Crude extract                  | <i>In vivo</i> (mice)   | [96]      |
| Mammary          | <i>Vitis vinifera</i>            | Grape             | Crude extract                  | <i>In vivo</i> (rats)   | [95]      |
| Skin cancer      | <i>Punica granatum</i> L.        | Pomegranate       | Crude extract                  | <i>In vivo</i> (mice)   | [92, 93]  |

**Table 2.** Anticarcinogenic effects of anthocyanin/anthocyanin-rich extract from different berry species under *in vivo* and *in vitro* conditions in different chronic diseases.

were associated with an increase in serum antioxidant capacity. At molecular level, Stoner et al. [90] also reported that the use of extracts showed a differential expression in 626 and 625 genes per 4807 and 17846 of preneoplastic esophagus and esophageal papilloma genes, respectively. These genes are involved in carbohydrate and lipid metabolism, cell death and proliferation, and inflammation. These results are an important approach to estimate the relation of anthocyanin gene expression and its influence on proteins associated with cell proliferation, apoptosis, angiogenesis, and esophageal carcinogenesis. Lala et al. [91] observed an anticarcinogenic effect of anthocyanins on colon cancer induced by

azoxymethane in a rat model. In that study, anthocyanin-rich extract from bilberry, chokeberry, and grapes significantly reduced azoxymethane-induced aberrant crypt foci and decreased cell proliferation and COX-2 gene expression [91]. Delphinidin and pomegranate extracts enriched with anthocyanins, and tannins showed an inhibition in skin cancer induced by UV-B or TPA (12-O tetradecanoylphorbol-13acetate) when applied to mouse skin [92, 93]. Here, delphinidin inhibited DNA damage mediated by UV-B radiation, and pomegranate modulated the mitogen-activated protein kinase (MAPK) and nuclear factor-kappa B (NF- $\kappa$ B) pathways. Similarly, studies on the protective effect and antioxidant mechanism of anthocyanin extract from blueberries were conducted using a liver injury induced by CC4 in mice—the effect of which increased lipid peroxidation and reduced liver cell viability [94]. The results indicate that anthocyanin extract effectively protected mice from CC4-induced liver injury by attenuation of lipid peroxidation. In mammary adenocarcinoma induced by dimethylbenzanthracene (DMBA) in rats, the antitumoral effect of grape juice was evaluated by Singletary et al. [95]. They demonstrated that the tumor mass was ultimately reduced by suppressing cell proliferation (**Table 2**). In general, the strong antioxidant capacity of berry species is attributed to their anthocyanin content, suggesting that it might offer potential chemopreventive properties, including the inhibition of gastric, leukemia, liver, and breast cancer cell proliferation, among others; however, the mechanism of action must be evaluated for each disease because apparently their mechanism of effects varies (inhibiting cell proliferation, activating different enzymatic activity, inducing or repressing gene expression, etc.) depending on the extract from each plant species.

## 5. Conclusions and future challenges

The potential use of anthocyanins from different plant species as natural compounds with a health benefit for humans opens a new trend for the prevention and alternative treatments of chronic diseases. Several reports have demonstrated that anthocyanins from berries could inhibit or decrease the growth of carcinogenic tumors by affecting cell proliferation, increasing or inhibiting enzymatic systems, and increasing expression of genes involved in cell protection. On the other hand, it is important to highlight that synthesis of anthocyanins in different tissues of plants species should be considered. In addition, the discovery and characterization of new regulatory elements of anthocyanin biosynthesis are crucial to understand and manipulate this pathway in breeding programs. Improving knowledge about increasing anthocyanin synthesis in crops of research and commercial interest, together with more animal and human model studies under *in vivo* conditions, is essential to generate better human anticarcinogenic or antichronic disease supplement products with chemopreventive effects from berries.

## Acknowledgements

We are very grateful to PIA-UFRO 16-0006 and DI 16-2013 projects from the Dirección de Investigación at the Universidad de La Frontera, Temuco, Chile.

## Author details

Daniela Peña-Sanhueza<sup>1</sup>, Claudio Inostroza-Blancheteau<sup>1, 2</sup>, Alejandra Ribera-Fonseca<sup>1, 3</sup> and Marjorie Reyes-Díaz<sup>1, 4\*</sup>

\*Address all correspondence to: [marjorie.reyes@ufrontera.cl](mailto:marjorie.reyes@ufrontera.cl)

1 Center of Plant, Soil Interaction and Natural Resources Biotechnology, Scientific and Technological Bioresource Nucleus (BIOREN), Universidad de La Frontera, Temuco, Chile

2 Núcleo de Investigación en Producción Alimentaria (NIPA-UCT), Escuela de Agronomía, Facultad de Recursos Naturales, Universidad Católica de Temuco, Temuco, Chile

3 Departamento de Producción Agropecuaria, Facultad de Ciencias Agropecuarias y Forestales, Universidad de La Frontera, Temuco, Chile

4 Departamento de Ciencias Químicas y Recursos Naturales, Facultad de Ingeniería y Ciencias, Universidad de La Frontera, Temuco, Chile

## References

- [1] Paredes-López O, Cervantes-Ceja ML, Vigna-Pérez M, Hernández-Pérez T. Berries: improving human health and healthy aging, and promoting quality. A review. *Plant Foods for Human Nutrition*. 2010;65:299–308.
- [2] Benvenuti S, Pellati F, Melegari M, Bertelli D. Polyphenols, anthocyanins, ascorbic acid, and radical scavenging activity of *Rubus*, *Ribes*, and *Aronia*. *Journal of the Science of Food and Agriculture*. 2004;69:164–169.
- [3] Céspedes C, El-Hafidi H, Pavon N, Alarcon J. Antioxidant and cardioprotective activities of phenolic extracts from fruits of Chilean blackberry *Aristotelia chilensis* (Elaeocarpaceae), Maqui. *Food Chemistry*. 2008;107:820–829.
- [4] Ruiz A, Gutierrez I, Mardones C, Vergara C, Herlitz E, Vega M, Dorau C, Winterhalter P, Von Baer D. Polyphenols and antioxidant activity of Calafate (*Berberis microphylla* G. Forst.) fruit and other native berries from southern Chile. *Journal of Agricultural and Food Chemistry*. 2010;58:6081–6089.
- [5] Jimenez S, Guevara R, Miranda R, Feregrino A, Torres I, Vazquez M. A review: functional properties and quality characteristics of bioactive compounds in berries: biochemistry, biotechnology, and genomics. *Food Research International*. 2012;54:1195–1207.
- [6] Koes RE, Quattrocchio F, Mol JNM. The flavonoid biosynthetic pathway in plants: function and evolution. *BioEssays*. 1994;16:123–132.
- [7] Boss P, Davies Ch, Robinson S. Expression of anthocyanin biosynthesis pathway genes in red and white grapes. *Plant Molecular Biology*. 1996;32: 565–569.

- [8] de Pascual-Teresa S, Moreno D, García-Viguera C. Flavonols and Anthocyanins in cardiovascular health: a review of current evidence. *International Journal of Molecular Sciences*. 2010;11:1679–1703.
- [9] Konczak I, Zhang W. Anthocyanins-more than natures colours. *Journal of Biomedicine and Biotechnology*. 2004;5:239–240.
- [10] Kong JM, Chia LS, Goh NK, Chia TF, Brouillard R. Analysis and biological activities of anthocyanins. *Phytochemistry*. 2003;64:923–933.
- [11] Soriano R, Pastore G. Evaluation of the effects of anthocyanins in type 2 diabetes. *Food Research International*. 2012;46:378–386.
- [12] Pazmino-Duran AE, Giusti MM, Wrolstad RE, Gloria BA. Anthocyanins from *Oxalis triangularis* as potential food colorants. *Journal of Agricultural and Food Chemistry*. 2001;75:211–216.
- [13] Castaneda-Ovando A, Pacheco-Hernandez M, Paez-Hernandez M, Rodriguez J, Galan-Vidal C. Chemical studies of anthocyanins: a review. *Food Chemistry*. 2008;113:859–871.
- [14] Nichenametla SN, Taruscio TG, Barney DL, Exon JH. A review of the effects and mechanisms of polyphenolics in cancer. *Critical Reviews in Food Science and Nutrition*. 2006;46:161–183.
- [15] Stintzing FC, Carle R. Functional properties of anthocyanins and betalains in plants, food, and in human nutrition. *Trends in Food Science and Technology*. 2004;15:19–38.
- [16] Szajdek A, Borowska EJ. Bioactive compounds and health promoting properties of berry fruits: review. *Plant Foods for Human Nutrition*. 2008;63:147–156.
- [17] Castrejón ADR, Eichholz I, Rohn S, Kroh LW, Huyskens-Keil S. Phenolic profile and antioxidant activity of highbush blueberry (*Vaccinium corymbosum* L.) during fruit maturation and ripening. *Food Chemistry*. 2008;109:564–572.
- [18] Connor AM, Luby JJ, Hancock JF, Berkheimer S, Hanson EJ. Changes in fruit antioxidant activity among blueberry cultivars during cold temperature storage. *Journal Agriculture and Food Chemistry*. 2002;50:893–898.
- [19] Diplock AT, Charleux J-L, Crozier-Willi G, Kok FJ, Rice-Evans C, Roberfroid M. Functional food science and defence against reactive oxidative species. *British Journal of Nutrition*. 1998;80:77–112.
- [20] Rice-Evans C, Miller NJ, Paganga G. Antioxidant properties of phenolic compounds. *Trends in Plant Science*. 1997;2:152–159.
- [21] Prior RL, Cao G. Antioxidant phytochemicals in fruits and vegetables: diet and health implications. *HortScience*. 2000;35:588–592.



- [22] Procházková D, Boušová I, Wilhelmov N. Antioxidant and prooxidant properties of flavonoids. Review. *Fitoterapia*. 2011;82:513–523.
- [23] Pieta PG. Flavonoids and antioxidants. *Journal of Natural Products*. 2000;63:1035–1042.
- [24] Nijveldt RJ, van Nood E, van Hoorn DEC, Boelens PG, van Norren K, van Leeuwen PAM. Flavonoids: a review of probable mechanisms of action and potential applications. *The American Journal of Clinical Nutrition*. 2001;74:418–25.
- [25] Moyer R, Hummer K, Finn C, Frei B, Wrolstad R. Anthocyanins, phenolics, and antioxidant capacity in diverse small fruits: *Vaccinium*, *Rubus*, and *Ribes*. *Journal of Agricultural and Food Chemistry*. 2002;50:519–525.
- [26] Pantelidis GE, Vasilakakis Manganaris GA, Diamantidis G. Antioxidant capacity, phenol, anthocyanin and ascorbic acid contents in raspberries, blackberries, red currants, gooseberries and Cornelian cherries. *Journal of Agricultural and Food Chemistry*. 2007;102:777–783.
- [27] Wada L, Ou B. Antioxidant activity and phenolic content of Oregon caneberries. *Journal of Agricultural and Food Chemistry*. 2002;50:3495–3500.
- [28] Prior R, Cao G, Martin A, Sofic E, McEwen J, O'Brien C, Lischner N, Ehlenfeldt M, Kalt W, Krewer G, Mainland M. Antioxidant capacity as influenced by total phenolic and anthocyanin content, maturity, and variety of *Vaccinium* species. *Journal of Agricultural and Food Chemistry*. 1998;46:2686–2693.
- [29] Ribera AE, Reyes-Diaz M, Alberdi M, Zuñiga GE, Mora ML. Antioxidant compounds in skin and pulp of fruits change among genotypes and maturity stages in highbush blueberry (*Vaccinium corymbosum* L.) grown in southern Chile. *Journal of Soil Science and Plant Nutrition*. 2010;10:509–536.
- [30] Rausher MD, Miler RE, Tiffin P. Patterns of evolutionary rate variation among genes on the anthocyanin biosynthetic pathway. *Molecular Biology and Evolution*. 1999;16:266–274.
- [31] Matus J, Loyola R, Vega A, Peña-Neira A, Bordeu E, Arce-Jhonson P, Alcald, J. Post-veraison sunlight exposure induces MYB-mediated transcriptional regulation of anthocyanin and flavonol synthesis in berry skins of *Vitis vinifera*. *Journal of Experimental Botany*. 2009;60:853–867.
- [32] Xi X, Zha Q, Jiang A, Tian Y. Impact of cluster thinning on transcriptional regulation of anthocyanin biosynthesis-related genes in “Summer Black” grapes. *Plant Physiology and Biochemistry*. 2016;104:180–187.
- [33] Li Y, Baldauf S, Lim EK, Bowles DJ. Phylogenetic analysis of the UDP-glycosyltransferase multigene family of *Arabidopsis thaliana*. *The Journal of Biological Chemistry*. 2001;276:4338–4343.

- [34] Hichri I, Barrieu F, Bogs J, Kappel Ch, Delrot S, Lauvergeat V. Recent advances in the transcriptional regulation of the biosynthetic pathway. Review paper. *Journal of Experimental Botany*. 2011;62:2465–2483.
- [35] Payne CT, Zhang F, Lloyd AM. GL3 encodes a bHLH protein that regulates trichome development in *Arabidopsis* through interaction with GL1 and TTG1. *Genetics*. 2000;156:1349–1362.
- [36] Baudry A, Caboche M, Lepiniec L. TT8 controls its own expression in a feedback regulation involving TTG1 and homologous MYB and bHLH factors, allowing a strong and cell-specific accumulation of flavonoids in *Arabidopsis thaliana*. *The Plant Journal*. 2006;46:768–779.
- [37] Matus J, Poupin M, Cañon P, Bordeu E, Alcalde J, Arce-Johnson P. Isolation of WDR and bHLH genes related to flavonoid synthesis in grapevine (*Vitis vinifera* L.). *Plant Molecular Biology*. 2010;72:607–620.
- [38] Paz-Ares J, Ghosal D, Wienand U, Peterson P, Saedler H. The regulatory c1 locus of *Zea mays* encodes a protein with homology to myb oncogene products and with structural similarities to transcriptional activators. *The EMBO Journal*. 1987;6:3553–3558.
- [39] Kobayashi S, Ishimaru M, Hiraoka K, Honda C. Myb-related genes of the Kyoho grape (*Vitis labruscana*) regulate anthocyanin biosynthesis. *Planta*. 2002;215:924–933.
- [40] Koes R, Verweij W, Quattrocchio F. Flavonoids: a colorful model for the regulation and evolution of biochemical pathways. *Trends in Plant Science*. 2005;10:236–242.
- [41] Matus J, Aquea F, Arce-Johnson P. Analysis of the grape MYB R2R3 subfamily reveals expanded wine quality-related clades and conserved gene structure organization across *Vitis* and *Arabidopsis* genomes. *BMC Plant Biology*. 2008;8:83.
- [42] Yang T, Perasso R, Baroin-Tourancheau A. MYB genes in ciliates: a common origin with the MYB proto-oncogene? *Protist*. 2003;154:229–238.
- [43] Lipsick JS. One billion years of MYB. *Oncogene*. 1996;13:223–235.
- [44] Weston K. Myb proteins in life, death and differentiation. *Current Opinion in Genetics & Development*. 1998;8:76–81.
- [45] Deluc L, Bogs J, Walker AR, Ferrier T, Decendit A, Merillon J-M, Robinson SP, Barrieu F. The transcription factor VvMYB5b contributes to the regulation of anthocyanin and proanthocyanidin biosynthesis in developing grape berries. *Plant Physiology*. 2008;147:2041–2053.
- [46] Terrier N, Torregrosa L, Ageorges A, Vialet S, Verries C, Cheynier V, Romieu C. Ectopic expression of VvMybPA2 promotes proanthocyanidin biosynthesis in *Vitis vinifera* L. and suggests additional targets in the pathway. *Plant Physiology*. 2009;149:1028–1041.
- [47] Stracke R, Werber M, Weisshaar B. The R2R3-MYB gene family in *Arabidopsis thaliana*. *Current Opinion in Plant Biology*. 2001;4:447–456.

- [48] Mehrtens F, Kranz H, Bednarek P, Wisshaar B. The Arabidopsis transcription factor MYB12 is a flavonol-specific regulator of phenylpropanoid biosynthesis. *Plant Physiology*. 2005;138:1083–1096.
- [49] Gonzalez A, Zhao M, Leavitt JM, Lloyd AM. Regulation of the anthocyanin biosynthetic pathway by the TTG1/bHLH/Myb transcriptional complex in Arabidopsis seedlings. *The Plant Journal*. 2007;53:814–827.
- [50] Aharoni A, Ric De Vos CH, Wein M, Sun Z, Greco R, Kroon A, Mol JNM, O'Connell AP. The strawberry FaMYB1 transcription factor suppresses anthocyanin and flavonol accumulation in transgenic tobacco. *The Plant Journal*. 2001;28:319–332.
- [51] Salvatierra A, Pimentel P, Moya-León MA, Herrera R. Increased accumulation of anthocyanins in *Fragaria chiloensis* fruits by transient suppression of FcMYB1 gene. *Phytochemistry*. 2013;90:25–36.
- [52] Takos AM, Jaffé FW, Jacob SR, Bogs J, Robinson SP, Walker AM. Light-induced expression of a MYB gene regulates anthocyanin biosynthesis in red apples. *Plant Physiology*. 2006;142:1216–1232.
- [53] Espley RV, Hellens RP, Putterill J, Stevenson DE, Kutty-Amma S, Allan AC. Red coloration in apple fruit is due to the activity of the MYB transcription factor, MdMYB10. *The Plant Journal*. 2007;49:414–427.
- [54] Ban Y, Honda C, Hatsuyama Y, Igarashi M, Bessho H, Moriguchi T. Isolation and functional analysis of a MYB transcription factor gene that is a key regulator for the development of red coloration in apple skin. *Plant Cell Physiology*. 2007;48:958–970.
- [55] Mathews H, Clendennen SK, Caldwell CG, Liu XL, Connors K, Matheis N, Schuster DK, Menasco DJ, Wagoner W, Lightner J, Wagner DR. Activation tagging in tomato identifies a transcriptional regulator of anthocyanin biosynthesis, modification, and transport. *The Plant Cell*. 2003;15:1689–1703.
- [56] Bogs J, Downey MO, Harvey JS, Ashton AR, Tanner GJ, Robinson SP. Proanthocyanidin synthesis and expression of genes encoding leucanthocyanidin reductase in developing grape berries and grapevine leaves. *Plant Physiology*. 2005;139:652–663.
- [57] Czemmel S, Stracke R, Weissahaarm B, Cordon N, Harris NN, Walker AR, Robinson SP, Bogs J. The grapevine R2R3-MYB transcription factor VvMYBF1 regulates flavonol synthesis in developing grape berries. *Plant Physiology*. 2009;151:1513–1530.
- [58] Lijavetzky D, Ruiz-García L, Cabezas JA, De Andrés MT, Bravo G, Ibañez A, Carreño J, Cabello F, Ibañez J, Martínez-Zapater JM. Molecular genetics of berry colour variation in table grape. *Molecular Genetics and Genomics*. 2006;276:427–435.
- [59] Walker AR, Lee E, Bogs J, McDavid DAJ, Thomas MR, Robinson SP. White grapes arose through the mutation of two similar and adjacent regulatory genes. *The Plant Journal*. 2007;49:772–785.
- [60] Lin-Wang K, Bolitho K, Grafton K, Kortstee A, Karunairetnam A, McGhie TK, Espley RV, Hellens RP, Allan AC. An R2R3 MYB transcription factor associated with regulation of the anthocyanin biosynthetic pathway in Rosaceae. *BMC Plant Biology*. 2010;10:50.

- [61] Toledo-Ortiz G, Huq E, Quail PH. The Arabidopsis basic/helix–loop–helix transcription factor family. *The Plant Cell*. 2003;15:1749–1770.
- [62] Li X, Duan X, Jiang H, Sun Y, Tang Y, Yuan Z, Guo J, Liang W, Chen L, Yin J, Ma H, Wang J, Zhang D. Genome-wide analysis of basic/helix–loop–helix transcription factor family in rice and Arabidopsis. *Plant Physiology*. 2006;141:1167–1184.
- [63] Ellenberger T, Fass D, Arnaud M, Harrison SC. Crystal structure of transcription factor E47: E-box recognition by a basic region helix–loop–helix dimer. *Genes & Development*. 1994;8:970–980.
- [64] Heim MA, Jakoby M, Werber M, Martin C, Weisshaar B, Bailey PC. The basic helix–loop–helix transcription factor family in plants: a genome-wide study of protein structure and functional diversity. *Molecular Biology and Evolution*. 2003;20:735–747.
- [65] Heisler MGB, Atkinson A, Bylstra YH, Walsh R, Smyth DR. SPATULA, a gene that controls development of carpel margin tissues in Arabidopsis, encodes a bHLH protein. *Development*. 2001;128:1089–1098.
- [66] Sorensen A-M, Krober S, Unte US, Huijser P, Dekker K, Saedler H. The Arabidopsis ABORTED MICROSPORES (AMS) gene encodes a MYC class transcription factor. *The Plant Journal*. 2003;33:413–423.
- [67] Abe H, Urao T, Ito T, Seki M, Shinozaki K, Yamaguchi-Shinozaki K. Arabidopsis AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling. *The Plant Cell*. 2003;15:63–78.
- [68] Li H, Sun J, Xu Y, Jiang H, Wu X, Li C. The bHLH-type transcription factor AtAIB positively regulates ABA response in Arabidopsis. *Plant Molecular Biology*. 2007;65:655–665.
- [69] Séguéla M, Briat JF, Vert G, Curie C. Cytokinins negatively regulate the root iron uptake machinery in Arabidopsis through a growth-dependent pathway. *The Plant Journal*. 2008;55:289–300.
- [70] Chandler VL, Radicella JP, Robbins TP, Chen J, Turks D. Two regulatory genes of the maize anthocyanin pathway are homologous: isolation of B utilizing R genomic sequences. *The Plant Cell*. 1989;1:1175–1183.
- [71] Ludwig SR, Habera LF, Dellaporta SL, Wessler SR. Lc, a member of the maize R gene family responsible for tissue-specific anthocyanin production, encodes a protein similar to transcription activators and contains the myc homology region. *Proceedings of the National Academy of Sciences USA*. 1989;86:7092–7096.
- [72] Goff SA, Klein TM, Roth BA, Fromm ME, Cone KC, Radicella JP, Chandler VP. Transactivation of anthocyanin biosynthetic genes following transfer of B regulatory genes into maize tissues. *EMBO Journal*. 1990;9:2517–2522.
- [73] Burr FA, Burr B, Scheffler BE, Blewitt M, Wienand U, Matz EC. The maize repressor-like gene intensifier shares homology with the rVb7 multigene family of transcription factors and exhibits missplicing. *The Plant Cell*. 1996;8:1249–1259.



- [74] Nesi N, Debeaujon I, Jond C, Pelletier G, Caboche M, Lepiniec L. The TT8 gene encodes a basic helix–loop–helix domain protein required for expression of DFR and BAN genes in *Arabidopsis* siliques. *The Plant Cell*. 2000;12:1863–1878.
- [75] Quattrocchio F, Wing JF, Leppen HTC, Mol JNM, Koes RE. Regulatory genes controlling anthocyanin pigmentation are functionally conserved among plant species and have distinct sets of target genes. *The Plant Cell*. 1993;5:1497–1512.
- [76] Quattrocchio F, Wing JF, van der Woude K, Mol JNM, Koes R. Analysis of bHLH and MYB domain proteins: Species-specific regulatory differences are caused by divergent evolution of target anthocyanin genes. *The Plant Journal*. 1998;13:475–488.
- [77] Smith TF, Gaitatzes C, Saxena K, Neer EJ. The WD repeat: a common architecture for diverse functions. *Trends in Biochemical Sciences*. 1999;24:181–185.
- [78] Van Nocker S, Ludwig P. The WD-repeat protein superfamily in *Arabidopsis*: conservation and divergence in structure and function. *BMC Genomics*. 2003;4:50.
- [79] Carey CC, Strahle JT, Selinger DA, Chandler VL. Mutations in the pale aleurone color1 regulatory gene of the *Zea mays* anthocyanin pathway have distinct phenotypes relative to the functionally similar TRANSPARENT TESTA GLABRA1 gene in *Arabidopsis thaliana*. *The Plant Cell*. 2004;16:450–464.
- [80] Walker AR, Davison PA, Bolognesi-Winfield AC, James CM, Srinivasan N, Blundell TL, Esch JJ, Marks MD, Gray JC. The TRANSPARENT TESTA GLABRA1 locus, which regulates trichome differentiation and anthocyanin biosynthesis in *Arabidopsis*, encodes a WD40 repeat protein. *The Plant Cell*. 1999;11:1337–1350.
- [81] de Vetten N, Quattrocchio F, Mol J, Koes R. The an11 locus controlling flower pigmentation in petunia encodes a novel WD-repeat protein conserved in yeast, plants, and animals. *Genes & Development*. 1997;11:1422–1434.
- [82] Sompornpailin K, Makita Y, Yamazaki M, Saito K. A WD-repeat-containing putative regulatory protein in anthocyanin biosynthesis in *Perilla frutescens*. *Plant Molecular Biology*. 2002;50:485–495.
- [83] Hanachi P, Kua SH, Asmah R, Motalleb G, Fauziah O. Cytotoxic effect of *Berberis vulgaris* fruit extract on the proliferation of human liver cancer cell line (HepG2) and its antioxidant properties. *International Journal of Cancer Research*. 2006;2:1–9.
- [84] Končić MZ, Kremer D, Karlović K, Kosalec I. Evaluation of antioxidant activities and phenolic content of *Berberis vulgaris* L. and *Berberis croatica* Horvat. *Food and Chemical Toxicology*. 2010;48:2176–2180.
- [85] Shin Y, Ryu JA, Liu RH, Nock JF, Watkins CB. Harvest maturity, storage temperature and relative humidity affect fruit quality, antioxidant contents and activity, and inhibition of cell proliferation of strawberry fruit. *Postharvest Biology and Technology*. 2008;49:201–209.
- [86] Chang YC, Huang HP, Hsu JD, Yang SF, Wang CJ. Hibiscus anthocyanins rich extract-induced apoptotic cell death in human promyelocytic leukemia cells. *Toxicology and Applied Pharmacology*. 2005;205:201–212.



- [87] Jia N, Xiong YL, Konga B, Liua Q, Xia X. Radical scavenging activity of black currant (*Ribes nigrum* L.) extract and its inhibitory effect on gastric cancer cell proliferation via induction of apoptosis. *Journal of Functional Foods*. 2012;4:382–390.
- [88] Huang H, Chang Y, Wu Ch, Hung Ch, Wang Ch. Anthocyanin-rich Mulberry extract inhibit the gastric cancer cell growth *in vitro* and xenograft mice by inducing signals of p38/p53 and c-jun. *Food Chemistry*. 2011;129:1703–1709.
- [89] Wang L, Stoner G. Anthocyanins and their role in cancer prevention. *Cancer Letters*. 2009;269:281–290.
- [90] Stoner G, Wang L-S, Seguin C, Rocha C, Stoner K, Chiu S, Kinghorn AD. Multiple berry types prevent N-nitrosomethylbenzylamine-induced esophageal cancer in rats. *Pharmacy Research*. 2010;27:1138–1145.
- [91] Lala G, Malik M, Zhao C, He J, Kwon Y, Giusti MM, Magnuson BA. Anthocyanin-rich extracts inhibit multiple biomarkers of colon cancer in rats. *Nutrition Cancer*. 2006;54:84–93.
- [92] Afaq F, Saleem M, Krueger CG, Reed JD, Mukhtar H. Anthocyanin- and hydrolysable tannin-rich pomegranate fruit extract modulates MAPK and NF-kappaB pathways and inhibits skin tumorigenesis in CD-1 mice. *International Journal of Cancer*. 2005;113:423–433.
- [93] Afaq F, Khan N, Syed DN, Mukhtar H. Oral feeding of pomegranate fruit extract inhibits early biomarkers UVB radiation induced carcinogenesis in SKH-1hairless mouse epidermis. *Photochemistry and Photobiology*. 2010;86:1318–1326.
- [94] Chen K, Sun H, Sun A, Lin Q, Wang Y, Tao X. Studies pf the protective effect and antioxidant mechanism of blueberry anthocyanins in a CC14-induced liver injury model mice. *Food and Agricultural Innumonoly*. 2012;23:352–362.
- [95] Singletary KW, Jung KJ, Giusti MM. Anthocyanin-rich grape extract blocks breast cell DNA damage. *Journal of Medical Food*. 2007;10:244–251.
- [96] Casto BC, Kresty LA, Kraly CL. Chemoprevention of oral cancer by black raspberries. *Anticancer Research*. 2002;22:4005–4015.